

Table 7. Resistance to Isoniazid with or without Rifampin Resistance in Reported TB Cases with No Previous TB by Origin: United States, 1993-1999

Year	Resistant to Isoniazid ^a						Resistant to Isoniazid and Rifampin ^a					
	Total Cases ^b		U. S.-born		Foreign-born ^c		Total Cases ^b		U. S.-born		Foreign-born ^c	
	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
1993	1,400	(8.4)	804	(6.8)	579	(12.4)	410	(2.5)	302	(2.6)	105	(2.3)
1994	1,355	(8.3)	709	(6.5)	632	(12.1)	352	(2.2)	238	(2.2)	109	(2.1)
1995	1,171	(7.3)	554	(5.4)	615	(11.0)	252	(1.6)	168	(1.6)	84	(1.5)
1996	1,134	(7.4)	492	(5.2)	638	(11.3)	205	(1.3)	103	(1.1)	101	(1.8)
1997	1,076	(7.5)	429	(5.0)	640	(11.3)	157	(1.1)	75	(0.9)	81	(1.4)
1998	1,012	(7.6)	366	(4.8)	644	(11.4)	127	(1.0)	52	(0.7)	74	(1.3)
1999	865	(7.2)	271	(4.1)	590	(11.1)	127	(1.1)	38	(0.6)	88	(1.7)

^aIsolates may be resistant to other drugs.

^bIncludes persons of unknown country of birth.

^cIncludes persons born outside the United States, American Samoa, the Federated States of Micronesia, Guam, the Republic of the Marshall Islands, Midway Island, the Commonwealth of the Northern Mariana Islands, Puerto Rico, the Republic of Palau, U.S. Minor Outlying Islands, U.S. Miscellaneous Pacific Islands, and the U.S. Virgin Islands.

Note: Data for all years updated through May 3, 2000.

More than 85% of all cases in each group had drug susceptibility test results reported for an initial isolate.

See Surveillance Slides #14 and #15.

Table 8. Resistance to Isoniazid with or without Rifampin Resistance in Reported TB Cases with Previous TB by Origin: United States, 1993-1999

Year	Resistant to Isoniazid ^a						Resistant to Isoniazid and Rifampin ^a					
	Total Cases ^b		U. S.-born		Foreign-born ^c		Total Cases ^b		U. S.-born		Foreign-born ^c	
	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
1993	164	(16.6)	85	(12.6)	76	(25.2)	75	(7.6)	30	(4.5)	45	(15.0)
1994	177	(17.1)	81	(11.7)	95	(28.1)	75	(7.3)	35	(5.1)	39	(11.6)
1995	168	(17.6)	77	(13.0)	91	(25.1)	70	(7.3)	28	(4.7)	42	(11.6)
1996	141	(16.4)	67	(12.0)	73	(24.2)	42	(4.9)	20	(3.6)	21	(7.0)
1997	108	(14.9)	35	(8.0)	73	(25.6)	42	(5.8)	11	(2.5)	31	(10.9)
1998	97	(12.8)	37	(7.5)	60	(22.6)	23	(3.0)	6	(1.2)	17	(6.4)
1999	80	(12.2)	27	(7.2)	51	(18.3)	25	(3.8)	6	(1.6)	19	(6.8)

^aIsolates may be resistant to other drugs.

^bIncludes persons of unknown country of birth.

^cIncludes persons born outside the United States, American Samoa, the Federated States of Micronesia, Guam, the Republic of the Marshall Islands, Midway Island, the Commonwealth of the Northern Mariana Islands, Puerto Rico, the Republic of Palau, U.S. Minor Outlying Islands, U.S. Miscellaneous Pacific Islands, and the U.S. Virgin Islands.

Note: Data for all years updated through May 3, 2000.

More than 85% of all cases in each group had drug susceptibility test results reported for an initial isolate.